



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

90. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Find the greatest number of inch balls that can be placed in a box 10 inches square and 5 inches deep.

91. Proposed by RAYMOND D. SMITH, Tiffin, Ohio.

A barn 20 feet square is standing in a pasture, and a horse is tied to one corner of it with a rope 50 feet long. Over how much land can he graze?

92. Proposed by J. A. CALDERHEAD, M. Sc., Professor of Mathematics in Curry University, Pittsburg, Pa.

What rate of income do I realize by purchasing United States 4% bonds at 105 if I sell them in six years at 104?

*** Solutions of these problems should be sent to B. F. Finkel, not later than April 10.

GEOMETRY.

88. Proposed by FREDERICK R. HONEY, Ph. B., Instructor in Mathematics in Trinity College, New Haven, Conn.

Prove that the volume of the frustum of a cone is equal to one-sixth of the altitude multiplied by the sum of the areas of the upper base, the lower base, and four times the area of the section midway between the upper and lower bases.

89. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Describe a circle tangent to three given circles. [From *Chauvenett's Geometry*, page 318, ex. 213.]

90. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va.

The bisectors of the angles of the opposite sides (produced) of an inscribed quadrilateral cut the sides at the angular points of a rhombus.

*** Solutions of these problems should be sent to B. F. Finkel, not later than April 10.

CALCULUS.

70. Proposed by J. OWEN MAHONEY, B. E., M. Sc., Graduate Fellow in Mathematics in Vanderbilt University, P. O., Lynnville, Tenn.

$$\text{Prove } \int_0^\infty \frac{\cos ax}{1+x^{2n}} dx = -i \frac{\pi}{n} \sum_{r=1}^n \omega^{2r-1} e^{ai} \omega^{2r-1}$$

where n is an integer, a is positive, and ω is $e^{i\pi/2n}$.

Is this correct? Forsyth gives, on page 41, of his *Theory of Functions*, the integral

$$\int_{-\infty}^{\infty} \frac{\cos ax dx}{1+x^{2n}} = -i \frac{\pi}{2l} \sum_{r=1}^n \omega^{2r-1} e^{ai} \omega^{2r-1}.$$